

Actiferol 15 mg

Sachets Food supplement. For dietary treatment to ensure proper supply of iron and in states of iron deficiency and anemia for infants and children.

Iron is a microelement required for proper development and functioning of the body. It contributes to proper development of cognitive functions in children. It contributes to the proper creation of hemoglobin and red cells which is really important for the growing body of a child. It helps in proper functioning of immunity system. It is an ingredient of hemoglobin which conditions transport of oxygen.



Actiferol Fe[®] contains unique form of iron put under micronization and emulgation. Thanks to micronization, i.e. great fragmentation, absorption of iron from iron pyrophosphate (III) is twice as high which is confirmed by researches. Iron contained in Actiferol Fe[®] is liberated in intestines, not in stomach. Thanks to that, there are no afflictions connected with digestive system.

Content:

| Nutritional value | 100 g | Recommended daily intake-1 sachet |
|--------------------------|--------------------|-----------------------------------|
| Energy | 1475 kJ (348 kcal) | 15 kJ (3 kcal) |
| Fat, of which: | 0 g | 0 g |
| Carbohydrates, of which: | 0 g | 0 g |
| • Sugar | 87 g | 0,9 g |
| Protein | 27 g | < 0,5 g |
| Salt | 0 g | 0 g |
| Iron | 0,63 g | 0,01 g |
| Carbohydrates, of which: | 1500 mg | 15 mg |

Ingredients: maltodextrin,, iron (ferric diphosphate (III) (ferric pyrophosphate), maltodextrin, emulsifier: lecithins.

How to use:

Infants and children to 12 years old: taken under doctor's supervision, depending on individual demand. Children above 12 years old: 1 sachet of iron 15 mg daily. In states of iron deficiency and/or anemia: taken under doctor's supervision, depending on individual demand.

Dissolve content of sachet in small amount of warm liquid (about ¼ of glass of water or breast milk) or add to food (one

spoon of porridge or soup). Consume directly after preparing.

Important information: Product taken under doctor's supervision. For dietary treatment. Product cannot be used parenterally. It isn't the only source of food. Do not use if there is hypersensitivity to any of the ingredients of the product. Daily intake of the substance cannot be reached only by change of standard diet. Varied diet and healthy lifestyle are important to stay healthy.

Packing:

• 30 sachets